

DTIC FILE COPY  
AD-A230 384

②

AD \_\_\_\_\_

REPORT NO 11-91

PROLONGED TREADMILL LOAD CARRIAGE:  
ACUTE INJURIES AND CHANGES IN FOOT ANTHROPOMETRY

U S ARMY RESEARCH INSTITUTE  
OF  
ENVIRONMENTAL MEDICINE  
Natick, Massachusetts

DTIC  
ELECTE  
JAN 02 1991  
S D CS D



Approved for public release distribution unlimited

UNITED STATES ARMY  
MEDICAL RESEARCH & DEVELOPMENT COMMAND

90 12 27 033

The findings in this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

The views, opinions and/or findings contained in this report are those of the authors and should not be interpreted as official Department of the Army position, policy, or decision, unless so designated by other official documentation.

Human subjects gave their free and informed voluntary consent before participating in this study. Investigators adhered to AR 70-25 and USAMRDC Regulation 70-25 on Use of Volunteers in Research.

Citations of commercial organizations and trade names in this report do not constitute an official Department of the Army endorsement or approval of the products or services of these organizations.

Approved for public release; distribution is unlimited.

#### DISPOSITION INSTRUCTIONS

Destroy this report when no longer needed.

Do not return to the originator.

PROLONGED TREADMILL LOAD CARRIAGE: ACUTE INJURIES AND CHANGES IN FOOT ANTHROPOMETRY

Katy L. Reynolds, MAJ, MC

SPC John Kaszuba

Robert P. Mello, M.A.

John F. Patton, Ph.D.

Occupational Health and Performance Directorate

US Army Research Institute of Environmental Medicine

Natick, Massachusetts 01760-5007

June, 1990

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
1a. REPORT SECURITY CLASSIFICATION N/A			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY N/A			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release - distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE N/A					
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION USARIEM		6b. OFFICE SYMBOL (If applicable) SGRD-UE-OPM		7a. NAME OF MONITORING ORGANIZATION US Army Medical Research and Development Command	
6c. ADDRESS (City, State, and ZIP Code) Natick, MA 01760-5007			7b. ADDRESS (City, State, and ZIP Code) Ft. Detrick Frederick, MD 21702-5012		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (If applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c. ADDRESS (City, State, and ZIP Code)			10. SOURCE OF FUNDING NUMBERS		
		PROGRAM ELEMENT NO. 62787A		PROJECT NO. 3E162787A879	TASK NO. BF
				WORK UNIT ACCESSION NO. 134	
11. TITLE (Include Security Classification) Prolonged Treadmill Load Carriage: Acute Injuries and Changes in Foot Anthropometry					
12. PERSONAL AUTHOR(S) Katy L. Reynolds, SPC John Kaszuba, Robert P. Mello, John F. Patton					
13a. TYPE OF REPORT Final		13b. TIME COVERED FROM May 88 TO May 89		14. DATE OF REPORT (Year, Month, Day) June 1990	
15. PAGE COUNT					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	Blisters, Foot anthropometry, Load carriage, Treadmill march		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The objectives of this study were to: 1) determine the distribution of injuries occurring during loaded treadmill marching, 2) determine if there are significant anthropometric changes that occur in the foot during this marching, and 3) identify variables (speed, load) that may affect these injuries and foot changes. Fifteen male subjects marched on a level treadmill at three speeds (3.96, 4.86, 5.76 km/hr), and with each of three loads (5.2, 31.5, 49.4 kg), in random order over a 7 week period. The march distance was 12 km and the Army's external frame pack system (Alice) was used. Aerobic power (maximum oxygen uptake) and body composition (body density by underwater weighing) were measured during the initial week. During the next 7 weeks foot anthropometric measurements were obtained before and after each treadmill march. A physician was present to record all injuries that occurred during each march. This study documented a large number of minor injuries (N=82) among physically fit soldiers marching on a treadmill at fast speeds and over a long distance with heavy loads. Significant pre to post foot anthropometric changes (p<.05) were observed in the ball of					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Katy L. Reynolds			22b. TELEPHONE (Include Area Code) (508) 651-4800		22c. OFFICE SYMBOL SGRD-UE-OPM

19. ABSTRACT (continued)

foot and instep circumferences with a tendency for these changes to occur with the heavier loads while ankle circumference changes occurred throughout most of the conditions. The results of this study indicate that: 1) load carriage injuries occurring during a prolonged treadmill march primarily involve the lower extremities and back, 2) foot blisters were the most common injury recorded, 3) the standard Army practice of using foot powder and undergoing frequent sock changes does not prevent blister formation during prolonged treadmill marches, 4) the most common types of back injury was upper back strain, and 5) transient rucksack palsy can occur when carrying a heavy load on a treadmill for an extended period of time, 6) marching on a treadmill over a prolonged period will significantly increase the circumference of the foot, and 7) treadmill load carriage march injuries are similar to actual field loaded road march injuries.

# CONTENTS

	PAGE
List of Figures and Tables	vi
Foreward	viii
Acknowledgements	ix
Executive Summary	1
Introduction	3
Methods	3
Results	7
Discussion	18
Conclusion	22
Recommendations	23
References	24
Appendixes	27

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification .....	
By .....	
Distribution / .....	
Availability Codes	
Dist	Avail and/or Special
A-1	



## LIST OF FIGURES AND TABLES

	PAGE
<b>FIGURES</b>	
1. Anthropometric foot measurements	6
2. Mean pre to post change in ball of foot breadth for each load carriage condition	8
3. Mean pre to post change in ball of foot circumference for each load carriage condition	9
4. Mean pre to post change in instep circumference for each load carriage condition	10
5. Mean pre to post change in heel breadth for each load carriage condition	11
6. Mean pre to post change in ankle circumference for each load carriage condition	12
7. Total injuries relative to load carriage weight	14
8. Total injuries relative to march rate	15

## **TABLES**

## **PAGE**

1. Load carriage conditions	4
2. Types of injuries for all load carriage conditions	13
3. Location of injuries for all load carriage conditions	16
4. Influence of load carriage conditions on foot injuries	16
5. Influence of load carriage conditions on foot blisters	17
6. Influence of load carriage conditions on upper back strain	17
7. Influence of load carriage conditions on shoulder injuries	18



## FOREWORD

Technology has changed many aspects of the US Army. Nevertheless, the Light Infantryman continues to rely on his own mobility in the battlefield. The soldier must transport his own equipment and be self-sustained for a period of several days until resupply is provided. Thus, it is important to understand the factors and problems related to the individual soldier's mobility, particularly over long distances and with heavy loads.

The Army recently created four new "light" infantry divisions. Loaded road march missions and training are common features in these divisions. The greater emphasis on loaded road march training has led to a resurgence of musculoskeletal injuries. These injuries are a major cost to the military in terms of lost manpower and health care cost (1). In the few studies which have systematically collected load carriage injury data, foot blisters appear to be the predominant injury (2,3,4). Other injuries have reported have included stress fractures (5), plantar fasciitis (2,3) and rucksack palsy (6).

In November 1988, the Surgeon General tasked the US Army Medical Research and Development Command to conduct a program of training injury research. While studying the physiology of load carriage, we availed ourselves of the opportunity to document load carriage-related injuries during prolonged walking on a motor driven treadmill. Since foot blisters are a common walking-related injury, this report also examines the foot morphometric changes that occur during loaded treadmill marching.

## **ACKNOWLEDGEMENTS**

The authors wish to acknowledge the technical expertise of Dr. Kenneth Parham (Natick Research, Development and Engineering Center) and the technical assistance of SPC Susan Covell and SGT Matthew Bovee. We also thank the subjects for their dedication and untiring efforts throughout this study.

## EXECUTIVE SUMMARY

The purposes of this study were to: 1) determine the distribution of injuries occurring during loaded treadmill marching, 2) determine if there are any significant foot anthropometric changes that occur during this marching, and 3) identify variables (speed, load) that may affect these changes. Fifteen male subjects marched on a level treadmill at three speeds (3.96, 4.86, 5.76 km/hr) and each of three loads (5.2, 31.5, 49.4 kg). The march distance was 12 km and the Army's external frame pack system (Alice) was used. Aerobic power (maximum oxygen uptake) and body composition (body density by underwater weighing) were measured. Foot anthropometric measurements were obtained before and after each treadmill march. A physician was present to record all injuries that occurred during each march. The results of this study indicate that 1) load carriage injuries during a prolonged treadmill march primarily involve the lower extremities and back, 2) foot blisters are the most common injury, 3) the standard Army practice of using foot powder and undergoing frequent sock changes does not prevent blister formation during prolonged treadmill marches, 4) the most common type of back injury is upper back strain, 5) transient rucksack palsy can occur when carrying a heavy load on a treadmill for an extended period of time, 6) Marching on a treadmill over a prolonged period will significantly increase the circumference of the foot, and 7) treadmill loadcarriage march injuries are similar to actual field loaded road march injuries. Recommendations include 1) the standard Army practice of preparing the feet prior to a road march needs to be systematically reevaluated and updated, 2) adequate planning is necessary before participating in a prolonged loaded march and unit leaders should consider several factors (footwear, load weight and distribution, etc.) during the preparation phase, 3) the sizing of footwear needs further evaluation to determine the relationship of these factors to the occurrence of foot blisters, and 4) anatomical and biomechanical factors play an important role in loaded marching and should be carefully considered in order to prevent injuries.

## INTRODUCTION

Load carriage is a vital part of military operations among the light infantry. Despite technical advances, the infantry soldier must in many cases depend on his own mobility to move himself and his equipment during combat. In certain field operations, an individual soldier may have to carry an amount of ammunition and equipment approaching his own body weight. This may lead to excessive fatigue and injuries as was experienced among the U.S. Army in Grenada (7). There is very little reliable data available on injuries occurring during prolonged load carriage marches. In a study conducted in Sweden (2), as many as 90% of the 114 infantry soldiers required medical attention for injuries encountered during a 20-26 km loaded roadmarch. In a recent US Army infantry exercise, Knapik et al, (3) reported that 24% of his subjects had one or more injuries while performing a maximal effort 20km road march and carrying a 46 kg load.

Foot blisters are the most common injuries reported (2,3,4) during load carriage marches. Cooper (4) related the incidence of blistering to increases in foot girth occurring during loaded road marching. However, there is very little information available on foot anthropometry and loaded road marching (8).

Other common injuries associated with load carriage include upper and lower back strain, metatarsalgia, plantar fasciitis, knee and ankle pain (2,3), stress fractures (5) and rucksack palsy (6). These injuries can debilitate the soldier and lead to lost mission effectiveness, training time, and ultimately his combat readiness.

The objectives of this study were three-fold, 1) to determine the distribution of injuries occurring during loaded treadmill marching, 2) to determine if there are significant anthropometric changes that occur during this marching, and 3) identify variables (speed, load) that may affect these injuries and foot changes.

## METHODS

**Subjects.** Fifteen male soldiers were recruited after completing either advanced individual training at Ft. Jackson or one-station unit training at Ft. Benning. All subjects were briefed about the study and signed a Volunteer Agreement Affidavit according to Army Regulations 70-25 and 40-38 prior to participation in the study. They then were evaluated by a physician

for medical clearance to participate as test subjects.

**Design.** During the initial week the subjects became acquainted with the laboratory procedures, and practiced walking on the treadmill at the various speeds and loads defined for the study. Body composition (body density by underwater weighing) and aerobic power (maximal oxygen uptake) were also measured during this period (9).

Over a 7 week period, the subjects performed 9 randomly ordered load carriage trials at 0% grade on a treadmill (Quinton Model 24-72). The loads and speeds are displayed in Table 1.

---

TABLE 1. Load carriage conditions

<u>Condition</u>	<u>Load(kg)</u>	<u>Speed(km/hr)</u>
1	5.2	3.96
2	5.2	4.86
3	5.2	5.76
4	31.5	3.96
5	31.5	4.86
6	31.5	5.76
7	49.4	3.96
8	49.4	4.86
9	49.4	5.76

---

The 5.2 kg load consisted of the standard battle dress uniform, standard black leather combat boots, and kevlar helmet. For the 31.5 kg and 49.4 kg loads, 13.6 and 31.5 kg, respectively, were carried in an all-purpose, lightweight, individual carrying equipment (ALICE) system. The rest of the weight (17.9 kg) consisted of 12.7 kg distributed on the equipment belt and the 5.2 kg as described above.

The trials were randomly arranged for each subject to minimize any training effects. Each subject performed no more than two trials per week, with each trial being separated by at least two days. The distance was fixed at 12 km for each march trial. A 10 min rest period was taken each hour. The total time for each trial (including rest periods) varied according to the speed: 3.96 km/hr (212 min), 4.86 km/hr (168 min), 5.76 km/hr (145 min).

The subjects were interviewed and examined by a physician at the beginning and end of each treadmill march to learn of any injuries that may have resulted from the march. The physician then recorded findings on an injury record form (Appendix). An injury was defined as any untoward event that occurred during testing which led to an impairment of body function or structure.

The subjects were given clean socks and foot powder was applied to their feet on a daily basis to minimize blistering. All subjects wore the standard plastic mesh ventilating insert normally supplied with the boots.

**Foot Anthropometric Measurements.** Five measurements were made on the bare right foot before and after the treadmill marches according to the procedures of White (10). The subject remained standing with a load in an erect stance on a platform. The feet were positioned slightly apart so that the subject's weight was evenly distributed. The Martin sliding caliper was utilized to determine the ball of foot (BOF) breadth and heel breadth. The first and fifth metatarsophalangeal joints were used as landmarks for measuring ball of foot breadth. The heel breadth measurements used the base of the Achilles tendon and the posterior aspects of the malleoli as landmarks.

The Gulich tape measure was used to measure the ball of foot (BOF), instep and ankle circumferences. The ball of foot circumference was determined by placing the tape over the first and fifth metatarsophalangeal joints. The instep circumference was measured by placing the tape measure at the highest point of the medial aspect of the arch of the foot and bringing it around the dorsum of the foot to the lateral aspect of the arch. The ankle circumference was determined by using the medial and lateral malleoli as the landmarks (Fig 1).

All measurements were performed twice, averaged, and recorded in millimeters.

**Statistical Analysis.** A one-way analysis of variance with repeated measures was used to determine pre to post changes of foot anthropometric measurements at each load condition. A two-way analysis of variance with repeated measures was also performed to find out if there were significant differences in foot girth pre to post test and if these differences were affected by load or speed. An alpha level of 0.05 was used to indicate statistical significance.




Figure 1. Foot anthropometric measurements.

## RESULTS

The subject descriptive characteristics were (mean  $\pm$  SD): age,  $21.1 \pm 1.0$  yrs; body mass,  $77.1 \pm 28$  kg; height,  $173.9 \pm 1.9$  cm; percent body fat,  $17.4 \pm 0.8$  %; and maximal oxygen uptake ( $\dot{V}O_{2\max}$ ),  $58.5 \pm 1.5$  ml $\cdot$ kg $^{-1}\cdot$ min $^{-1}$ . When compared to other Army populations, the mean ( $\pm$ SD) for percent body fat was similar (11) but the  $\dot{V}O_{2\max}$  was slightly higher than the average soldier completing advanced training (12).

Figures 2-6 describe foot anthropometric changes at the various speeds and loads. No pre to post changes were noted with ball of foot breadth (diagonal) for any condition. Significant changes were noted in the ball of foot circumference with combinations of 1) a light load at a fast speed, 2) a medium load at a medium speed, and 3) a heavy load at a slow speed. Instep circumference changes were significant with the medium and heavy loads at both slow, medium and fast speeds. Heel breadth measurement changes were significant with the light load at both the slow and fast speeds and with the medium load at a medium pace. Ankle circumferences showed a significant increase during all load carriage conditions except for the medium load at a medium speed.

The two-way analysis of variance revealed that there were significant differences ( $p < .05$ ) pre to post tests in all foot anthropometric measurements except for ball of foot breadth (diagonal). However, the repeated measures revealed no significant pre to post differences by load or speed.

Injuries were recorded during each load carriage condition as shown in Table 2. There were a total of 82 injuries reported and friction blisters and upper back strain accounted for more than 50% of the total.






Figure 2. Mean pre and post values for ball of foot breadth for each load carriage condition. (\*  $p < .05$ )




Figure 3. Mean pre and post values for ball of foot circumference for each load carriage condition. (\*  $p < .05$ )




Figure 4. Mean pre and post values for instep circumference for each load carriage condition. (\*  $p < .05$ )




Figure 5. Mean pre and post values for heel breadth for each load carriage condition. (\*  $p < .05$ )




Figure 6. Mean pre and post values for ankle circumference for each load carriage condition. (\*  $p < .05$ )

---

TABLE 2. Types of injuries for all load carriage conditions.

<u>TYPE</u>	<u>NUMBER</u>	<u>PERCENT</u>
Friction Blisters (feet)	24	29.4%
Upper back strain	19	23.3%
Hot Spots (feet)	10	12.2%
Shoulder numbness	2	2.5%
Shoulder tendonitis	4	4.9%
Metatarsal Pain (feet)	4	4.9%
Abrasions (shoulder/abdomen)	3	3.6%
Plantar Fasciitis (feet)	3	3.6%
Tendonitis (knee)	3	3.6%
Toenail Injury	3	3.6%
Groin Strain	2	2.5%
Low Back Strain	1	1.2%
Hip Pain	1	1.2%
Ankle Sprain	1	1.2%
Foot Contusions	1	1.2%
Metatarsal Fracture (feet)	1	1.2%
<b>TOTAL</b>	<b>82</b>	<b>100.0</b>

---

Figure 7 displays the percentage of total injuries relative to load carriage weight. Sixty-five percent (53/82) of the injuries occurred with the heaviest load.

Figure 8 shows the percentage of total injuries with respect to march rate. Forty-five percent (37/82) of the injuries occurred at the fastest speed.

Table 3 shows the location of injuries for all conditions. More than 60% of the total injuries involved the lower extremities with the feet being the most predominant body part injured. The back was the second most common injury site.




Figure 7. Total injuries relative to load carriage weight.




Figure 8. Total injuries relative to march rate.



TABLE 3. Location of injuries for all load carriage conditions.

<u>LOCATION</u>	<u>NUMBER</u>	<u>PERCENT</u>
Feet	46	56.1%
Back	20	24.5%
Shoulders	8	9.8%
Knees	3	3.6%
Groin	2	2.4%
Hips	1	1.2%
Ankle	1	1.2%
Abdomen	1	1.2%
<b>TOTAL</b>	<b>82</b>	<b>100.0</b>

The influence of the load carriage conditions on foot injuries is shown in Table 4. A greater percentage of the foot injuries occurred with the heaviest load at the medium and fast speeds. Carrying a medium load at both medium and fast speeds also led to frequent injuries.

TABLE 4. Influence of load carriage conditions on foot injuries.

<u>CONDITIONS</u>	<u>NO.OF INJURIES</u>	<u>PERCENT</u>
1. 5.2kg, 3.96 km/hr	0	0.0%
2. 5.2kg, 4.86 km/hr	1	2.2%
3. 5.2kg, 5.76 km/hr	3	6.5%
4. 31.5kg, 3.96 km/hr	3	6.5%
5. 31.5kg, 4.86 km/hr	8	17.4%
6. 31.5kg, 5.76 km/hr	6	13.1%
7. 49.4kg, 3.96 km/hr	0	0.0%
8. 49.4kg, 4.86 km/hr	10	21.7%
9. 49.4kg, 5.76 km/hr	15	32.6%
<b>TOTAL</b>	<b>46</b>	<b>100.0</b>

The effects of the load carriage condition on foot blisters are summarized in Table 5. A

majority of the friction blisters were observed with the heavy load at both medium and fast speeds.

---

TABLE 5. Influence of load carriage conditions on foot blisters.

<u>CONDITIONS</u>	<u>NO. OF INJURIES</u>	<u>PERCENT</u>
1. 5.2 kg, 3.96 km/hr	0	0.0%
2. 5.2 kg, 4.86 km/hr	1	4.2%
3. 5.2 kg, 5.76 km/hr	2	8.3%
4. 31.5 kg, 3.96 km/hr	2	8.3%
5. 31.5 kg, 4.86 km/hr	3	12.5%
6. 31.5 kg, 5.76 km/hr	1	4.2%
7. 49.4 kg, 3.96 km/hr	0	0.0%
8. 49.4 kg, 4.86 km/hr	7	29.2%
9. 49.4 kg, 5.76 km/hr	8	33.3%
<b>TOTAL</b>	<b>24</b>	<b>100.0</b>

---

The effect of the various conditions on upper back strain is shown in Table 6. The frequency of upper back strain was highest with the heaviest load during fast, medium and slow speeds.

---

TABLE 6. Influence of load carriage conditions on upper back strain.

<u>CONDITIONS</u>	<u>NO. OF INJURIES</u>	<u>PERCENT</u>
1. 5.2 kg, 3.96 km/hr	0	0.0%
2. 5.2 kg, 4.86 km/hr	0	0.0%
3. 5.2 kg, 5.76 km/hr	0	0.0%
4. 31.5 kg, 3.96 km/hr	0	0.0%
5. 31.5 kg, 4.86 km/hr	2	10.5%
6. 31.5 kg, 5.76 km/hr	2	10.5%
7. 49.4 kg, 3.96 km/hr	6	31.6%
8. 49.4 kg, 4.86 km/hr	6	31.6%
9. 49.4 kg, 5.76 km/hr	3	15.8%
<b>TOTAL</b>	<b>19</b>	<b>100.0</b>

---

Table 7 shows the influence of the load carriage conditions on the frequency of shoulder injuries. The majority of shoulder injuries occurred with the heaviest load at both slow and fast speeds. Two cases of upper extremity numbness were reported at the medium and heavy loads at medium speeds. The symptoms resolved when the packs were removed.

Table 7. Influence of load carriage conditions on shoulder injuries.

<u>CONDITIONS</u>	<u>NO. OF INJURIES</u>	<u>PERCENT</u>
1. 5.2 kg, 3.96 km/hr	0	0.0%
2. 5.2 kg, 4.86 km/hr	0	0.0%
3. 5.2 kg, 5.76 km/hr	0	0.0%
4. 31.5 kg, 3.96 km/hr	0	0.0%
5. 31.5 kg, 4.86 km/hr	1	12.5%
6. 31.5 kg, 5.76 km/hr	0	0.0%
7. 49.4 kg, 3.96 km/hr	2	25.0%
8. 49.4 kg, 4.86 km/hr	1	12.5%
9. 49.4 kg, 5.76 km/hr	4	50.0%
<b>TOTAL</b>	<b>8</b>	<b>100.0</b>

## DISCUSSION

The purpose of this study was to document the types and incidence rates of injuries occurring during a prolonged treadmill march, carrying various loads at different speeds. The authors also wanted to examine the effects of certain variables (speed, weight) on changes in foot anthropometry.

The injuries predominantly involved the lower extremities and back. Foot blisters were the most common injury recorded and this has been observed in field loaded roadmarches (2,3,4). Blisters can debilitate the soldier and lead to complications of infection and cellulitis (13). Friction blisters are due to frictional shearing forces that are applied to the epidermis of the skin (13,14). Moist feet can influence the time required for blistering by effecting the coefficient of friction for the skin (13). The standard Army practice of using foot powder and undergoing frequent sock changes are measures that have been utilized to prevent blisters. Other preventive measures such as wearing an inner polypropylene or nylon sock have also

been reported to keep the feet dry by wicking away foot moisture (15,16). Reynolds, et al (17) reported that antiperspirants may be a helpful measure in reducing the frequency of blisters. Perhaps using special products to keep the feet dry would have resulted in a lower incidence of blisters in the present study.

The second most common site of injury in this study was the back (20%) which was higher than found in other studies (2,3). Differences could be attributed to the small sample size and the fact that the subjects carried heavier loads. Results showed that the highest number of back injuries occurred with the heaviest load. It has been shown that during treadmill walking, the rucksack is not in synchrony with the movements of the trunk muscles. The muscles must compensate for this asynchrony and over a period of time this leads to fatigue and injury (18). Also, trunk angle and vertical forces increase as the load is increased which results in more stress on the back muscles (19).

The most common type of injury involving the back was upper back strain. Electromyographic studies (20) reveal that the activity in the upper back muscles is higher when the pack is placed high on the back as was the case in this study. If these muscles were not conditioned for loadcarriage then fatigue and injury might be expected to occur.

The shoulder was the third most common injury site with two cases presenting as a transient rucksack palsy. Other authors have reported this injury during load carriage marches and the shoulder girdle and elbow flexors are the most common sites affected. Reported symptoms include paresthesias, numbness and weakness secondary to compression of the upper trunk of the brachial plexus (6,21). Factors contributing to this disorder include using rucksacks without a frame or hipbelt, and carrying heavy packs (6). The subjects in this study used frames and hip belts. However, the two cases of rucksack palsy did occur with the heavier loads. Symptoms decreased when the pack was adjusted and completely resolved when the pack was removed.

In the present study, we documented one stress fracture by xray and four cases of metatarsal pain. It was difficult to determine whether the stress fracture was related to the marching, to off duty activities (basketball), or to accumulative stresses from both activities. Stress fractures and metatarsal pain have been associated with loaded marching especially in troops undergoing initial military training (22,23). Knapik, et al (3) also reported metatarsal pain during a 20 km loaded road march conducted with light infantry soldiers. Several factors shown to be associated with foot injuries are low fitness level (23,24), excess body weight (24), long march distances (25), and type of training activities (26). In the present study, a large percentage of the foot injuries were noted with the heavier loads. Kinoshita (27) found that heavier loads led to foot rotation that exposed the distal end of the metatarsals to

mechanical stresses for a longer period of time. The author recommended that altering stride length and wearing flexible boots might decrease the forces of stress in this area. Cushion insoles have also been recommended to decrease these stresses (15) but other sources report contradictory findings (28). This study did not examine this issue.

Foot anthropometric measurements were performed in this study to determine if foot dimensions change during loaded marching. Freedman, et al (8) reported that the dimensions of the feet did not change significantly during prolonged marching with weights for 13.7 miles/day for 32 days. However, the subjects were tested in Army dress shoes and carried 20 lb weights in their hands. In contrast, Cooper (4) noted an increase in foot girth as high as 9 mm in 49 % of soldiers participating in a 56km night march while carrying a rifle and water bottle. In the present study, significant pre to post march changes were noted with heel breadth measurements and ball of foot, instep and ankle circumferences. However, the maximum changes were not as high as in Cooper's study.

The significant heel breadth changes did not show a clear relationship to march speed or load. Any changes in pre to post measurements were probably related to changes in gait during the different conditions.

Ankle circumference changes occurred in all but one condition. There was a tendency for the significant changes in ball of foot and instep circumferences to occur with the heavier loads. However, more extensive analysis revealed that these changes were not significant. Therefore, the circumference increases could have been related to the marching and not to the load weight. Foot venous capacitance increases have been noted during weightbearing exercise. Armstrong and Kenney (29) reported significant foot venous capacitance increases with strain gauge plethysmography before and after an 8 week running program.

Foot anthropometric changes were not significantly affected by speed in this study. However, it would be interesting to see the effect of march speed on foot girth over longer distances.

In summary, this study documented a large number of minor injuries among physically fit soldiers marching on a treadmill at various speeds and over a long distance with heavy loads. Also, there was a tendency for certain foot anthropometric changes to occur with the heavier loads and the association of load and speed to foot girth changes during marching needs further evaluation. There are still many unanswered questions about the nature and causes of injuries specifically related to loaded road marches. Additional laboratory and field research studies need to be conducted so that preventive strategies can be developed to minimize injuries. This would have a beneficial effect on both unit strength and combat readiness.

## **CONCLUSIONS**

1. Load carriage injuries occurring during a prolonged treadmill march primarily involve the lower extremities and back.
2. Foot blisters were the most common injury recorded.
3. The standard Army practice of using foot powder and undergoing frequent sock changes does not prevent blister formation during prolonged treadmill marches.
4. The most common type of back injury was upper back strain.
5. Transient rucksack palsy can occur when carrying a heavy load on a treadmill for an extended period of time.
6. Marching on a treadmill over a prolonged period will significantly increase the circumference of the foot.
7. Treadmill load carriage march injuries are similar to actual field loaded road march injuries.

## **RECOMMENDATIONS**

1. The standard Army practice of preparing the feet prior to a road march needs to be systematically reevaluated and updated.
2. Adequate planning is necessary before participating in a prolonged loaded march and unit leaders should consider several factors (footwear, load weight and distribution, etc.) during the preparation phase.
3. The sizing of footwear for the soldier needs further evaluation to determine the relationship of these factors to the occurrence of foot blisters.
4. Anatomical and biomechanical factors play an important role in loaded marching and should be carefully considered in order to prevent injuries.

## REFERENCES

1. Tomlinson JP, Ledmar WM, Jackson JD. Risk of injury in soldiers. *Military Medicine* 152: 60-64, 1987.
2. Dalen A, Nilsson J, Thorstensson A: Factors influencing a prolonged foot march. Stockholm, Sweden (Karolinska Institute), FOA Report No.C 50601-H6, 1978.
3. Knapik J, Reynolds K, Staab J, Vogel J. Injuries resulting from a strenuous road march. (unpublished).
4. Cooper DS. Research into foot lesions among Canadian field forces. 13th Commonwealth Defense Conference on Operational Clothing and Combat Equipment, Paper CDA II, 1981.
5. Brudvig TGS, Gudger TD, Obermeyer L. Stress fractures in 295 trainees: a one-year study of incidence as related to age, sex, and race. *Military Medicine* 148:666-667, 1983.
6. Besson RJ, Belcher VW, Franklin RJ. Rucksack paralysis with and without rucksack frames. *Military Medicine* 152:372-375, July 1990.
7. Dubick JM, Fullerton TD. Soldier overloading in Grenada. *Military Review* pg 38-47, Jan 1987.
8. Freedman A., Kirkpatrick CM, Huntington EC. Survey of foot measurements and the proper fit of Army shoes. Armored Medical Research Laboratory Technical Report No. T13/45, Dec 1945.
9. Patton J, Kaszuba J, Mello RP, Reynolds KL. Physiological and perceptual responses to prolonged treadmill loadcarriage. US Army Research Institute of Environmental Medicine Technical Report No. T11-90, February 1990.
10. White RM. Comparative anthropometry of the foot. U.S. Army Natick Research, Development and Engineering Center Technical Report No. T10/83, Dec 1982.
11. Vogel JA, Kirkpatrick JW, Fitzgerald PI, Hodgdon JA, Harmon EA. Derivation of anthropometry based body fat equations for the Army's weight control program. US Army Research Institute of Environmental Medicine Technical Report No. 17/88, May 1988.



12. Vogel JA, Patton JF Mello RP, Daniels WL. An analysis of aerobic capacity in a large United States population. *J Appl Physiol* 60:494-600, 1986.
13. Akers WM. Sulzberger on friction blistering. *Int J Derm* 16:369-372, June 1977.
14. Naylor, PFD. Experimental friction blisters. *British J Dermatology* 67:327-342, 1955.
15. Blaber PE, Egan KM. The multi-component boot system. *Infantry March/April*:14-17, 1989.
16. Puffer JC. Blisters. *Sports Medicine Digest*. Vol. 11 No. 12:7, 1989.
17. Reynolds KL, Darrigrand A, Jackson R, Roberts D, Hamlet M. Effects of pedal antiperspirants on sweat accumulation. *Med Sci Sports Exerc* 22:S781, April 1990.
18. Pierrynowski MR, Norman RW, Winter DA. Mechanical energy analysis of the human during load carriage on a treadmill. *Ergonomics* 24:1-14, 1981.
19. Hale Cj, Coleman FR, Karpovich PV. Trunk inclination in carrying low and high packs of various weights. U.S. Army Quartermaster Research and Development Center, Lawrence, Ma, Report No. 216, July 1953.
20. Bobet J, Norman RW. Effects of load placement on back muscle activity in load carriage. *Eur J Appl Physiol* 53:71-75, 1984.
21. Wilson WJ. Brachial plexus palsy in basic trainees. *Military Medicine* 152:519-522, 1988.
22. Hurlinger CCW, and Tyler WL. March fracture. Report of 313 cases. *Bull. U.S. Army Med. Dept.* 69:72-80, 1944.
23. Gardner I, Dziados JE, Jones BJ, Brundage JF, Harris JM, Sullivan R, Gill P. Prevention of lower extremity stress fractures: a controlled study of a shock absorbant insole. *Am J Pub Health* 78:1563-1567, Dec 1988.
24. Gilbert RS and Johnson, HA. Stress fractures in military recruits. *Military Medicine* 131:716-721, 1966.
25. Jones BH, Manikowski R, Harris JH, Dziados J, Norton S, Ewart T, Vogel JA. Incidence of and risk factors for injury an illness among male and female army basic trainees. U.S.

Army Research Institute of Environmental Medicine Technical Report No. T19/88, 1988.

26. Scully TJ, Besterman G. Stress fractures-a preventable training injury. *Military Medicine* 147:285-286, 1982.

27. Kinoshita H. Effects of different load and carry systems on selected biomechanical parameters describing gait. *Ergonomics* 28:1347-1362, 1985.

28. Bansel CK. Wear test for U.S. Army trainee testing of boot inserts, memo dtd 1 December 1986.

29. Armstrong CG, Kenney WL. Functional effects of weight bearing exercise on the veins of the foot. *Med Sci Sports Exer* 19:S81, 1987.

APPENDIX  
DATA FORM USED TO RECORD INJURIES

INJURY DATA ENTRY FORM

DATE \_\_/\_\_/\_\_

SUBJECT NO. \_\_\_\_\_

STUDY TITLE \_\_\_\_\_

NAME \_\_\_\_\_

LAST

FIRST

MI

INVESTIGATOR \_\_\_\_\_

RACE \_\_\_\_\_ SEX- F \_\_ YES \_\_ NO M \_\_ YES \_\_ NO

SSN# \_\_\_\_\_ - \_\_\_\_\_ - \_\_\_\_\_

TREADMILL- PRE [\_\_] POST [\_\_] FOLLOWUP [\_\_]

MARCH TIME \_\_\_\_\_ MIN.

MARCH PACE \_\_\_\_\_ KM/HR

RUCKSACK TYPE- ALICE YES ☐ NO ☐

STANDARD HIPBELT USED? YES ☐ NO ☐

LBE TYPE STANDARD ☐ VEST ☐ OTHER ☐

RUCKSACK WEIGHT \_\_\_\_\_ KG (TOTAL INCLUD. LBE + HELMET)

DID AN INJURY OCCUR?


YES ☐ NO ☐

IF YES, THEN COMPLETE THE FOLLOWING:

PLEASE SPECIFY DATE OF ACTUAL OCCURRENCE OF INJURY IF IT DIFFERS FROM THAT ON  
PAGE ONE.

DATE \_\_\_\_/\_\_\_\_/\_\_\_\_

LOCATION- 1. HEAD 2. NECK 3. SHOULDERS 4. CHEST 5. ARMS 6. HANDS 7. UPPER  
(PLEASE) BACK 8. MID BACK 9. LOWER BACK 10. SACRUM 11. PELVIS 12. HIPS  
(CIRCLE) 13. THIGHS 14. LEGS 15. ANKLES 16. FEET 17. OTHER \_\_\_\_\_



BODY SIDE OF INJURY -RIGHT \_\_\_\_YES \_\_\_\_NO LEFT \_\_\_\_YES \_\_\_\_NO

SIGNS/SYMPTOMS YES ☐ NO ☐

IF YES, WHAT WERE THEY?

1. TENDERNESS ☐ YES ☐ NO

2. LIMITED ROM OF JOINT ☐ YES ☐ NO

3. INFLAMMED JOINT (WARMTH OR ERYTHEMA OR SWELLING)  
☐ YES ☐ NO

4. DECREASED SENSATION BY PINPRICK ☐ YES ☐ NO

5. TINGLING ☐ YES ☐ NO

6. MUSCLE SPASM ☐ YES ☐ NO

7. WEAKNESS ☐ YES ☐ NO

8. SKIN ABRASION ☐ YES ☐ NO

9. CONTACT RASH ☐ YES ☐ NO

10. SKIN INFECTION \_\_\_ YES \_\_\_ NO

11. BLISTER \_\_\_ YES \_\_\_ NO

12. BLACK TOENAIL \_\_\_ YES \_\_\_ NO

13. TORN TOENAIL \_\_\_ YES \_\_\_ NO

14. ATHLETES FEET \_\_\_ YES \_\_\_ NO

15. OTHER \_\_\_\_\_

DIAGNOSIS \_\_\_\_\_  
\_\_\_\_\_

XRAY \_\_\_ YES \_\_\_ NO IF YES, THEN WHAT TYPE? \_\_\_ PLAIN XRAY \_\_\_\_\_ RESULTS



\_\_\_\_\_ BONE SCAN \_\_\_\_\_ RESULTS

\_\_\_\_\_ OTHER (SPECIFY)

\_\_\_\_\_ RESULTS

TREATMENT 1. \_\_\_\_\_

2. \_\_\_\_\_

3. \_\_\_\_\_

4. \_\_\_\_\_

DISPOSITION

LIMITATIONS YES ☐ NO ☐ IF YES, THEN ☐ UPPER BODY HOW LONG? \_\_\_\_\_

☐ MID BODY HOW LONG ? \_\_\_\_\_

☐ LOWER BODY HOW LONG? \_\_\_\_\_

OTHER \_\_\_\_\_

QUARTERS YES ☐ NO ☐ IF YES, HOW LONG \_\_\_\_\_ HRS

HOSPITAL YES ☐ NO ☐ IF YES , HOW LONG \_\_\_\_\_ DAYS

OTHER \_\_\_\_\_

WHEN DID SUBJECT RETURN TO THE STUDY? \_\_\_\_\_

FULL PARTICIPATION \_\_\_ YES \_\_\_ NO

IF NO, THEN PLACE DATE WHEN THEY ACTUALLY RETURNED TO FULL PARTICIPATION.

\_\_\_\_/\_\_\_\_/\_\_\_\_

IF NEVER THEN STATE REASON \_\_\_\_\_

\_\_\_\_\_.

## DISTRIBUTION LIST

### 4 Copies to:

Defense Technical Information Center  
ATTN: DTIC-DDA  
Alexandria, VA 22304-6145

### 2 Copies to:

Commander  
U.S. Army Medical Research and Development Command  
ATTN: SGRD-OP  
Fort Detrick  
Frederick, MD 21702-5012

Commander  
U.S. Army Medical Research and Development Command  
ATTN: SGRD-PLE  
Fort Detrick  
Frederick, MD 21702-5012

Commander  
U.S. Army Medical Research and Development Command  
ATTN: SGRD-PLC  
Fort Detrick  
Frederick, MD 21702-5012

### 1 Copy to:

Commandant  
Academy of Health Sciences, U.S. Army  
ATTN: AHS-COM  
Fort Sam Houston, TX 78234-6100

Stimson Library  
Academy of Health Sciences, U.S. Army  
ATTN: Chief Librarian  
Bldg. 2840, Room 106  
Fort Sam Houston, TX 78234-6100

Director, Biological Sciences Division  
Office of Naval Research - Code 141  
800 N. Quincy Street  
Arlington, VA 22217

Commanding Officer  
Naval Medical Research and Development Command  
NMC-NMR/ Bldg. 1  
Bethesda, MD 20814-5044

Office of Undersecretary of Defense for Acquisition  
ATTN: Director, Defense Research and Engineering  
Deputy Undersecretary for Research & Advanced Technology  
(Environmental and Life Sciences)  
Pentagon, Rm. 3D129  
Washington D.C. 20301-3100

Dean  
School of Medicine  
Uniformed Services University Of The Health Sciences  
4301 Jones Bridge Road  
Bethesda, MD 20814-4799

2 Copies to:

Commander  
U.S. Army Medical Research Institute of Chemical Defense  
Aberdeen Proving Ground, MD 21010-5425

Commander  
U.S. Army Chemical Research, Development and Engineering Center  
Aberdeen Proving Ground, MD 21010-5423

Commandant  
U.S. Army Chemical School  
Fort McClellan, AL 36205-5020

Commander  
U.S. Air Force School of Aerospace Medicine  
Brooks Air Force Base, TX 78235-5000

Commander  
Naval Health Research Center  
P.O. Box 85122  
San Diego, CA 92138-9174

Commander  
U.S. Army Biomedical Research and Development Laboratory  
Fort Detrick  
Frederick, MD 21702-5010

Commander  
U.S. Army Medical Materiel Development Activity  
Fort Detrick  
Frederick, MD 21702-5009

U.S. Army Scientific Liaison Officer to DCIEM  
(U.S. Army Medical R&D Command)  
1133 Sheppard Avenue W.  
P.O. Box 2000  
Downsview, Ontario  
CANADA M3M 3B9

1 Copy:

Commandant  
Walter Reed Army Institute of Research  
Walter Reed Army Medical Center  
ATTN: SGRD-UWZ-C (Director for Research Management)  
Washington D.C. 20307-5100

Commander  
U.S. Army Environmental Hygiene Agency  
Aberdeen Proving Ground, MD 21010-5422

Director  
Army Physical Fitness Research Institute  
Box 469  
U. S. Army War College  
Carlisle Barracks, PA 17013

Commandant  
Army Physical Fitness Center  
U. S. Army Soldier Support Institute  
Fort Benjamin Harrison, IN 46216

Director  
Army Personnel Research Establishment  
c/o RAE, Farnborough  
Hants, GU14 6TD  
United Kingdom

HQDA  
SGPS-FP  
5109 Leesburg Pike  
Falls Church, VA 22041-3258